



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Invasive plants threaten the least mobile butterflies in Switzerland

Gallien, Laure ; Altermatt, Florian ; Wiemers, Martin ; Schweiger, Oliver ; Zimmermann, Niklaus E

Abstract: Aim: Biological invasions are today the second-largest global threat for biodiversity. Once introduced, exotic plant species can modify ecosystem composition, structure and dynamics, eventually driving native species to local extinction. Among the groups of organisms, most likely to be directly affected by exotic invasive plants are herbivorous insects, such as butterflies, which strongly depend on plants throughout their life cycle. However, it remains unclear whether invasive plants have a negative or a positive effect on butterfly diversity at a landscape scale. Location: Switzerland. Methods: Using an extensive inventory (393 sites across Switzerland) of both butterfly and invasive plants, we explore the impact of 31 invasive black listed plant species on local butterfly richness. We further identify each butterfly species' response to invasive plants (i.e. positive, neutral or negative) and analyse the functional and phylogenetic characteristics of these different groups of species. Results: Our results indicate that butterfly richness negatively correlates with an increase in invasive plant richness. When studying the individual response of each butterfly species to the number of invasive plants, we found that no single butterfly is profiting from invasive plant species, while 28 butterfly species (24%) suffer from the presence of invasive plants. We further show that the species negatively affected are on average less mobile than the unaffected species and that they are phylogenetically clustered. Main conclusions: Our results present evidences of the influence of invasive species on other trophic levels and interaction networks. We further highlight that a lack of management efforts for mitigating invasive plant impacts threatens specific sections of the functional and phylogenetic diversity of butterflies.

DOI: <https://doi.org/10.1111/ddi.12513>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-139401>

Journal Article

Accepted Version

Originally published at:

Gallien, Laure; Altermatt, Florian; Wiemers, Martin; Schweiger, Oliver; Zimmermann, Niklaus E (2017). Invasive plants threaten the least mobile butterflies in Switzerland. *Diversity and Distributions*, 23(2):185-195.

DOI: <https://doi.org/10.1111/ddi.12513>

Invasive plants threaten the least mobile butterflies in Switzerland

Laure Gallien¹, Florian Altermatt^{2,3}, Martin Wiemers⁴, Oliver Schweiger⁴ & Niklaus E. Zimmermann¹

¹ Swiss Federal Institute for Forest, Snow and Landscape Research WSL, CH-8903 Birmensdorf, Switzerland

²Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department of Aquatic Ecology, Überlandstrasse 133, CH-8600 Dübendorf, Switzerland.

³Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstr. 190, CH-8057 Zürich, Switzerland.

⁴UFZ – Helmholtz Centre for Environmental Research, Theodor-Lieser-Str. 4, 06120 Halle, Germany

E-mails: laure.gallien@gmail.com; florian.altermatt@eawag.ch; martin.wiemers@ufz.de; oliver.schweiger@ufz.de; niklaus.zimmermann@wsl.ch

Corresponding author: Laure Gallien

Running head: Plant invaders threaten butterflies

Article type: Biodiversity Research and Reviews

Word count: Abstract (263), Main text (4575).

Number of references: 61

ABSTRACT

Aim. Biological invasions are today the second-largest global threat for biodiversity. Once introduced, exotic plant species can modify ecosystem composition, structure and dynamics, eventually driving native species to local extinction. Among the groups of organisms most likely to be directly affected by exotic invasive plants are herbivorous insects, such as butterflies, which strongly depend on plants throughout their life cycle. However, it remains unclear whether invasive plants have a negative or a positive effect on butterfly diversity at a landscape scale.

Location. Switzerland.

Methods. Using an extensive inventory (393 sites across Switzerland) of both butterfly and invasive plants we explore the impact of 31 invasive black listed plant species on local butterfly richness. We further identify each butterfly species' response to invasive plants (i.e. positive, neutral, or negative), and analyse the functional and phylogenetic characteristics of these different groups of species.

Results. Our results indicate that butterfly richness negatively correlates with an increase in invasive plant richness. When studying the individual response of each butterfly species to the number of invasive plants, we found that no single butterfly is profiting from invasive plant species, while 28 butterfly species (24%) suffer from the presence of invasive plants. We further show that the species negatively affected are on average less mobile than the unaffected species, and that they are phylogenetically clustered.

Main conclusions. Our results present evidences of the influence of invasive species on other trophic levels and interaction networks. We further highlight that a lack of management efforts for mitigating invasive plant impacts threatens specific sections of the functional and phylogenetic diversity of butterflies.

46 Keywords: biological invasions, black listed species, functional diversity, invasion impacts,
47 land-use, phylogenetic diversity.

48

INTRODUCTION

Numerous invasive alien plant species can modify ecosystem properties, and replace native plant species, sometimes leading to their local extinction (Sax & Gaines, 2008; Morales & Traveset, 2009; Vilà *et al.*, 2011). Although some native species may suffer from invasions by exotic plants, such as native plants being out-competed or pollinators losing their feeding resources (e.g. Morales *et al.*, 2013; Vilcinskas *et al.*, 2013; Ignace & Chesson, 2014; Villéger *et al.*, 2015; see Schweiger *et al.*, 2010 for a review), some native species may profit from the presence of invasive plants, such as native herbivores gaining a new feeding resource (e.g. Foster & Robinson, 2007; Masters & Emery, 2015).

Among the groups of organisms likely to be directly affected by invasive plant species are butterflies, which strongly depend on plants throughout their life cycle (caterpillars feed on plants as herbivores, while adults are using plants for nectaring; e.g. Altermatt & Pearse, 2011). Invasive plant species can indeed affect butterflies' life cycles either in a positive or negative manner. On the one hand, butterflies may benefit from invasive plants if they provide additional or better quality food resources (e.g. more or higher quality of nectar; Graves *et al.*, 2003; Jahner *et al.*, 2011; Pearse & Altermatt, 2013). On the other hand, butterflies may suffer from plant invasions if these replace beneficial native plant partners, attract predators, or are toxic for the butterflies that feed on them (e.g. Tallamy & Shropshire, 2009; Davis & Cipollini, 2014, see also Bezemer, *et al.* 2014; van Hengstum *et al.*, 2014; Litt *et al.*, 2014 for similar examples on other arthropods).

Although interactions between invasive plants and native butterflies is receiving more attention at the local scale (e.g. Forister & Wilson, 2013; Pearse & Altermatt, 2013), broad scale patterns and general trends are currently understudied (Gallien & Carboni, 2016). At the landscape scale, it has for instance been shown that butterfly richness is affected by climate, landscape structure, and level of urbanization (Warren *et al.*, 2001; Forister *et al.*, 2010;

Concepción *et al.*, 2015, 2016). However, it remains unknown whether invasive plant richness also influences butterfly richness, and if so, whether this has a positive or a negative effect overall.

Similarly, when considering native butterfly species individually, it is likely that not all species will respond to plant invasions in the same manner. Indeed, it has been shown that narrow-ranged, diet specialist and dispersal limited butterfly species are most sensitive to global change (Warren *et al.*, 2001; Brook *et al.*, 2008). Species with small ranges have difficulties to maintain their populations viable when part of their range is threatened (e.g. Payne & Finnegan, 2007; Ohlemüller *et al.*, 2008), species with highly specialized diets cannot easily cope with novel ecosystems (e.g. Davies *et al.*, 2004), and dispersal limitations as well as long generation times are known to slow down a species' capacity to respond to local disturbances that can be either of biotic or abiotic nature (Weed *et al.*, 2013). Thus in the context of butterfly sensitivity to invasive plants, we can expect that very mobile species – which can more easily fly to suitable hosts – are less affected than less mobile species. Similarly, narrow ranged and specialist butterflies – which strongly depend on local resources – may be more affected by numerous invasive plants than wide ranging and generalist species. Many more important butterfly traits related to their susceptibility to invasive plant species may exist but often they are difficult to measure (e.g. color matching with host plants, attraction of natural enemies, adult nectar provider). However, if such features show phylogenetic signal (Münkemüller *et al.*, 2015) then butterflies suffering from plant invasions can be expected to be phylogenetically clustered.

Here we provide a first assessment of the influence of invasive plant richness on native butterfly richness and functional groups at the landscape scale. To do so, we take advantage of a large inventory effort on butterfly species, native and invasive black listed plant species in Switzerland, in which 393 transects were monitored (Fig. S1.1). Combining the analysis of

these surveys with information on butterfly species functional traits (such as mobility capacities, range size, or diet specificity) and phylogenetic position enables us to explore the influence of invasive plants on butterfly richness, and to identify the functional and phylogenetic characteristic of the most impacted butterflies.

Specifically, we provide a first assessment of whether and how native butterfly richness is impacted by invasive plant species richness along environmental gradients. Then, we identify those butterfly species that are significantly favoured or suppressed in areas with an increased number of invasive plant species. We further test for functional and phylogenetic differences between suppressed and non-suppressed butterflies. Finally, we provide an estimate of the geographic locations where butterflies are more likely to suffer from the presence of invasive plants, which may thus become priority areas for biodiversity protection.

METHODS

The Monitoring Scheme

Switzerland has a surface area of about 41,000 km² and encompasses a large diversity of environmental gradients since about 50% of its area is mountainous (elevations ranging from 190 m to 4634 m a.s.l.). To obtain a representative sample of the butterfly and plant diversity along these extended gradients, we used the Swiss Biodiversity Monitoring program dataset (BDM; BDM Coordination Office 2011, 2014). BDM monitors the biodiversity of Switzerland since 2001 and is composed of 393 sites of 1x1 km size that are regularly distributed over Switzerland (Fig. S1.1). Within each of these BDM sites, vascular plants and butterflies were surveyed between 2007 and 2011 using standardized methods (i.e. transects of 2.5 km along paths and roads within the 1 km² plots). Depending on elevation the sites were sampled once or twice for plants, and four to seven times for butterflies: annually

between April 21st and September 21st, with a time interval of at least 14 days (Altermatt *et al.*, 2008). A total of 1916 (native and exotic) plant species, and 187 butterfly species were identified (Fig. S1.2).

The black list of invasive exotic plant species was obtained from the Swiss National Centre of Plant Data and Information (www.infoflora.ch), and contains 56 highly invasive and dominant plant species exotic to Switzerland (*sensu* Richardson *et al.*, 2000) which show both a high spread potential, and a demonstrated negative impact on native biodiversity, human health and/or economy. For the following analyses, we selected in this list the species that were terrestrial and present in the BDM plant inventory (see list of the invaders in Appendix S1; black listed invader richness ranging from 0 to 15 species per site).

Functional Traits and Phylogeny

We used a set of functional traits describing the butterflies' range size (as a proxy for habitat specialization), larval diet specificity, number of generation per year, and morphological traits related to dispersal. As a coarse measure of the butterflies' European distribution range size we used the CLIMBER dataset (which covers all Europe and has a resolution of 50 x 50 km; Schweiger *et al.*, 2014). Diet breadth of the larval stages was based on previously published field-observed interactions between food plant species and Lepidoptera (data from Ebert, 1991-2005; Altermatt, 2010). Diet breadth of the larval stages was classified as monophagous (only one food plant species), strongly oligophagous (>1 food plant species, but only one food plant genus), oligophagous (>1 genus but only one food plant family), or polyphagous (>1 food plant family). We used information on the number of generations per year (after Ebert, 1991-2005), describing each species' voltinism. Finally, we measured the thorax width and forewing length on life-sized photographs of all species, in order to calculate wing-load ratios (i.e. ratio of thorax width to forewing length). Higher wing-load ratios are usually associated with higher dispersal abilities (Dudley & Srygley,

1994; Turlure *et al.*, 2010; but see Sekar, 2012).

The butterfly phylogeny is based on a molecular phylogenetic analysis of the concatenated mitochondrial gene *cytochrome c oxidase I* (1532 bp) and the nuclear gene *elongation factor I alpha* (1725 bp) of 85% of European butterfly species including all but two Swiss butterfly species (a total of 425 sequences). We used the Maximum Likelihood method based on a General Time Reversible model (Wiemers & Schweiger, unpublished), starting the heuristic search with trees obtained by applying the Neighbour-Joining method to the matrix of pairwise distances estimated using the Maximum Composite Likelihood approach. A discrete Gamma distribution was used to model evolutionary rate differences among sites. The phylogenetic tree with the highest log likelihood was chosen. This analysis was conducted with MEGA5 (Tamura *et al.*, 2011).

Environmental variables

To explore the relationships between plant and butterfly richness along environmental gradients we considered a large set of variables encompassing: topo-climate, habitat type and habitat diversity at different spatial resolutions, as well as the nearest distance to each habitat type (see Table S1.1 for a detailed list of variables). The topo-climatic variables included: annual sum of growing degree-days, mean annual temperature, mean annual potential evapotranspiration, mean annual moisture index, annual precipitation sum, number of frost days during the growing season, slope, aspect and elevation range per 1 km² pixel.

Statistical analyses

As a first step, we reduced the number of environmental variables by means of an ordination approach that has the advantage of providing uncorrelated axes of variation between sites in the subsequent regression analyses. We applied a principal component analysis (PCA) and used the first 6 PCA axes as representative of typical environmental

gradients. These axes accumulated to explain 76% of the inter-site differences (Fig. S1.3). The first PCA axis notably represented low temperatures, high proportion of forest cover, and long distances to urban and agricultural areas (Axis 1). The second PCA axis represented high habitat diversity and high proportion of urban areas (Axis 2). The third axis was correlated to high proportion of wetlands (Axis 3), while the fourth axis was related to high precipitation (Axis 4), the fifth axis to the number of frost days (Axis 5), and the sixth axis to the amount of solar radiation (Axis 6; Fig. S1.3). It can be noted that the proportion of urban cover, a variable important for both plants and butterflies, was well represented by the PCA axes as the 6 PCA axes together predict 81% of the urban cover variance in our dataset.

We aimed at estimating whether butterfly richness was affected by invasive plant richness, in addition to environmental variables and native plant richness. But since both native and invasive plant richness are also influenced by environmental variables, we adopted an approach avoiding spurious correlations between butterfly and invasive plant richness due to common correlations to environmental variables (for instance if both depend on temperature, they are likely to appear correlated). To do so, we implemented three analysis steps: First, we modelled native plant richness as a function of environmental variables (i.e. 6 PCA axes). Second, we modelled invasive plant richness with the environment and the residuals from the native plant richness model as predictors (i.e. the influence of the native plant richness that was not due to the environment). Third, we modelled butterfly richness as a function of the environmental variables, the residuals from the native plant richness model, the residuals from the invasive plant richness model (i.e. the influence of the native and invasive plant richness that were not due to the environment), and the interaction terms between the residuals from the invasive plant model and the environmental variables. For all models we allowed environmental variables to have both linear and quadratic terms.

For each regression we fitted a generalized linear model (GLM; using Poisson or quasi-

Poisson distributions). Full initial models were optimized by means of a stepwise backward-forward variable selection based on AIC scores optimization (the model with the lowest AIC score was retained; note that we used quasi-AIC for the quasi-Poisson models as suggested by Bolker 2014) with the aim to retain only the statistically relevant variables. All final model residuals were checked to comply with the normality assumption. The richness of invasive plant species was thus identified as being correlated to the overall richness of butterfly species when retained in the final set of variables, with an estimated coefficient being significantly different from 0. The explanatory power of the GLMs was evaluated by means of the regression R^2 (based on the proportion of deviance explained and corrected by the number of degrees of freedom), and variable importance was estimated using a 99-fold randomization procedure for each variable separately (Strobl *et al.*, 2007). The principle behind these randomizations was to mimic the absence of the variable in the model while maintaining the original degrees of freedom. The importance is then estimated as the average difference in prediction accuracy (i.e. regression R^2) with and without permuting the target variable. Note that we did not find spatial autocorrelation in the residuals of any of the fitted models (see mantel correlograms in Fig. S2.1).

To test whether individual butterfly species were particularly (positively or negatively) affected by local richness of exotic species, we then built logistic regression models for each butterfly species separately. In these models, the presence and absence of the species was explained by means of both the pre-selected environmental variables (i.e. 6 PCA axes) and the residuals of the native and invasive plant richness models (i.e. the influences of the native and invasive plant richness that were not due to the environment). We again applied a stepwise backward-forward variable selection procedure based on AIC scores (starting from a full model). For each butterfly species, the invasive plant species richness was considered to have a significant influence if it was retained in the stepwise-optimized model and had an estimated

coefficient significantly different from 0 (with a p-value corrected for multiple testing across each butterfly species; Benjamini & Yekutieli, 2001). Note that for statistical robustness, we analysed only those butterfly species that had at least 30 occurrences in our dataset (115 species in total, i.e. 62% of all butterfly species in the inventory).

It can be noted that all mentioned regressions rely on the assumption that at the landscape scale invasive plant species may influence butterfly species occurrence and richness, but are not affected by butterflies in return. This assumption is driven notably by the repeated evidence of an effect of specific invasive plants on native butterfly species (e.g. Tallamy & Shropshire, 2009; Jahner *et al.*, 2011; Davis & Cipollini, 2014), but none (to the best of our knowledge) about the landscape scale effects of native butterflies on invasive plants. The latter may be found though at the local scale.

Then, we classified the butterfly species according to their response to invasive plant richness into three groups: (i) those that were positively correlated (i.e. eventually profiting from invasive plant richness), (ii) those that were negatively correlated (i.e. eventually suffering from invasive plants richness), and (iii) those that did not reveal correlation with invasive plant richness. We then explored whether these species groups differed in their functional and phylogenetic characteristics. Specifically, we used ANOVA to test for functional differences in: (i) species' range size in Switzerland (as estimated by the number of sites in which they occur), (ii) species' range size in Europe, (iii) species' larval diet specificity, (iv) species' voltinism, and (v) species' wing-load ratio. The patterns of phylogenetic relatedness among species belonging to each of these groups were tested by means of the mean phylogenetic distance between all pairs of species (MPD; Webb *et al.*, 2002). To quantify whether the species within the groups were more clumped (clustered) or spread (overdispersed) across the phylogeny than expected by random grouping, we applied a null model randomizing the species' position on the tips of the phylogeny (999 repetitions),

and we calculated the standardized effect size (SES), hereafter called MPD_{SES} . MPD_{SES} can vary between 0 (completely clustered species) and 1 (completely overdispersed species), where 0.05 and 0.95 significance thresholds were applied.

Finally, in order to visualize the areas where the butterflies profiting or suffering from exotic species are located we mapped both the prediction (from the individual models generated in the second step) and observation of these species. To map the individual model predictions we transformed the continuous habitat suitability predictions into binary prediction with a threshold maximizing the proportion of correctly predicted presences (i.e. model sensitivity) and the proportion of correctly predicted absences (i.e. model specificity).

All statistical analyses were performed in R version 3.1.2 (R Development Core Team, 2014) and the packages: *ade4* (Dray & Dufour, 2007), *MASS* (Venables & Ripley, 2002), *bbmle* (Bolker & R Development Core Team, 2014), *ecodist* (Goslee & Urban, 2007), *PresenceAbsence* (Freeman & Moisen, 2008).

RESULTS

Plant and butterfly richness along the gradients

The selected set of environmental variables successfully predicted plant and butterfly richness, with explained deviances of $R^2=0.44$ for native plant richness, $R^2=0.65$ for invasive plant richness, and $R^2=0.75$ for butterfly richness. These three groups all responded negatively to the number of frost days during the growing season (PCA axis 5, Fig. 1; see Table S2.1 for the full list of estimated parameters of each model), but they responded differently to all other variables (Fig. 1).

The two most important variables for explaining butterfly richness in Switzerland were *environment-independent native plant richness* (40% with a positive coefficient; hereafter

called *native plant richness*) and the annual number of frost days (Axis 5, 26%; Fig. 2). The *environment-independent number of invasive plants* (hereafter called *invasive plants richness*) nonetheless improved the butterfly model significantly ($p\text{-value}<0.001$) and contributed to 4.1% of the explained deviance. The richness of invasive plant species revealed an overall negative effect on butterfly richness ($p\text{-value}<0.001$), and significant interactions with: axis 1 (temperature, forest cover, distance to urban and agricultural area), axis 3 (wetland cover), and axis 5 (annual number of frost days) as shown in Figure 2 and supplementary Table S2.1.

Characterizing butterflies that suffer from invasive plants

Among the 115 most abundant butterfly species modelled, we detected 28 species (24%; 16 genera) whose occurrence probability significantly decreased when invasive plant richness increases (Fig. 3b; see the full list of species in Appendix S2), but not a single species was positively affected. These butterfly species negatively associated with invasive plant richness had significantly smaller wing-load ratios than butterflies that were not affected by invasive plant species richness ($p\text{-value}<0.001$; Fig. 3a). All other tested traits showed no significant differences between butterfly categories: species range sizes in Switzerland ($p\text{-value}=0.60$) and in Europe ($p\text{-value}=0.82$), larval diet specificity ($p\text{-value}=0.06$), and average number of recorded generations per year ($p\text{-value}=0.16$). The phylogenetic distance between butterflies that are negatively related to invasive plant richness was significantly lower than expected by chance ($\text{MPD}_{\text{SES}}=0.016$). Note that we could not consider the two burnet moth species (Zygaenidae family; both negatively related to invasive plant richness) in this phylogenetic estimate because they were not included in the phylogeny.

The butterfly species that are negatively related to invasive plant richness were particularly present (both in number and as proportion of the local butterfly richness) in the

mountainous regions (i.e. north-western Switzerland in the Jura mountains, and southern Switzerland in the Swiss Alps; Fig. 4).

DISCUSSION

From the analysis of co-occurrence patterns between native butterflies and invasive plant species we explored the potential influence of invasive plant richness on butterfly diversity. Our results suggest that butterfly richness is negatively affected by invasive plant richness in Switzerland. Butterflies particularly sensitive to invasive plants seem to be the least mobile species, and phylogenetically clustered.

Invasive plant richness negatively correlates with butterfly richness

Invasive plant and butterfly species richness did not respond in the same way to the environment, with invasive plants preferring warmer and more urbanized sites compared to butterflies (Fig. 1, Axis 1). In fact, most invasive plants tend to originate from warm regions (Pyšek *et al.*, 2003; Walther *et al.*, 2007; Hulme, 2009) and are usually more abundant in urban and agricultural areas due to the high levels of disturbance in these areas and the high, human-induced propagule pressure (Chytrý *et al.*, 2008; Cabra Rivas *et al.*, 2015; Gallien *et al.*, 2015). Additionally, invasive plant richness was a negative and significant predictor of the overall butterfly species richness (even after removing the effect of the environment and native plant richness on invasive species richness).

We also found that the negative influence of invasive plant richness was specifically important in cold sites close to forested areas, in sites of low wetland cover, and in sites where the annual number of frost days is particularly high (see interactions in Fig. 2). These results indicate that in unsuitable conditions for most of the butterfly species, invasive plants may additionally decrease the local resources for the butterflies, for instance by replacing

beneficial native plant partners, by attracting predators, or by being toxic for the butterflies (e.g. Tallamy & Shropshire, 2009, Davis & Cipollini, 2014). However, in sites with a high proportion of wetland cover the invasive species richness had locally a seemingly positive effect on butterfly richness. These sites, however, are generally only suitable for a small set of highly specialized butterflies (see Axis 3 in Fig. 1), which are generally rare. Yet, we could not confirm these specific positive effects of invasive plants on butterfly richness at the butterfly species level, since our analyses were limited to the more common butterflies, while wetland habitat specialists are usually rare.

Furthermore, we found that the measure of invasive plant richness is a better indicator of butterfly richness than other variables that are sometimes used, such as the level of habitat diversity, or the proportion of urban and wetland cover (e.g. Concepción *et al.*, 2015; see Fig. 2). The explanatory power of invasive plant richness can, nonetheless, represent either a direct link to the butterfly physiology, with exotic species providing toxic resources for larvae and/or adult butterflies, or a missing covariate in the model, such as the frequency of disturbance in the sampled sites. It is essentially impossible to completely rule out the possible effect of missing covariates in empirical studies. However, we have used many covariates that are usually good proxies for disturbances (e.g. proportion of urban and agricultural fields, as well as habitat diversity). Also, the explanatory power of the model is high ($R^2=0.75$), and we found no spatial autocorrelation in the residuals (see Fig. S2.1). Therefore, it seems unlikely that an important covariate with spatial structure is missing.

It should be noted that we only had information available on the richness and not on specific abundances of invasive species per site (such as relative abundance or cover proportion). Substituting richness with specific dominance information in our analysis would likely reinforce our observed patterns (as improving the data quality should increase the explanatory power). Such information would be particularly useful to identify priority targets

for conservation measures (e.g. if one invader was highly dominant, it would greatly influence the presence and distribution of butterflies and should thus become a target for invasive management plans).

Least mobile butterflies are most affected by invasive plants

Not all butterfly species responded in the same way to invasive plants. Specifically, 24% of the analysed butterfly species were less likely to occur in sites with a high number of invasive plants. These vulnerable butterflies are significantly less mobile than the butterflies that are insensitive to the presence of invasive plant richness. This suggests that more mobile species are less affected than less mobile species because the former might more easily reach suitable host plants and escape sites affected by invasive plants (see also Warren *et al.*, 2001). The higher sensitivity of dispersal-limited butterflies to invasive plants furthermore suggests that the invasive plants do not only use local resources that are not used by the native plant species, but replaces native plant species on which the butterflies depend.

Butterflies' range size, specialization and voltinism do not seem to be significant drivers of their sensitivity to invasive plants. Nonetheless, since the sensitivity of butterfly species to invasive plant richness is a characteristic that is clustered in the butterfly phylogeny, additional functional traits (that show phylogenetic signals) may also be affected by invasive plants (such as male mate-searching behaviour or female oviposition behaviour; Pavoine *et al.*, 2014). Even though it may be difficult to quantitatively analyse many important butterfly traits that can be related to their susceptibility to invasive plant species, the position of butterfly species in the phylogeny may be used as a first indication for their general sensitivity. The phylogenetic clustering of sensitive butterfly species has important further implications, as it indicates that some phylogenetic clades – such as the genera *Polyommatus*

and *Argynnis* – are specifically sensitive to increasing numbers of invasive plants. As a consequence, increased introduction and invasion of exotic plant species can lead to a decrease in the phylogenetic diversity of local butterfly assemblages.

Although we did not test for specific pairwise interactions between individual invasive plant and butterfly species due to our study design, our results can be used as a first screening to identify particularly vulnerable butterfly species that would ideally be studied in a more specific pairwise plant-butterfly interaction context. Additionally, our screening approach can be used to target the butterflies most vulnerable to invasive plant richness. These species should be prioritized in conservation planning. However, we point to the fact that the already rare and endangered butterflies, which are likely also very vulnerable to invasive plants, could not be assessed in this study due to data scarcity.

The detected impact of invasive plants on both butterfly functional characteristics and phylogenetic diversity might lead to functional and phylogenetic homogenization of butterfly assemblages with increasing amounts of exotic plant species. Functional homogenization of butterfly communities has been already documented across multiple functional groups (Eskildsen *et al.*, 2015) and was found as a consequence of land use change (Ekroos *et al.*, 2010). Similar results were also suggested for bird species assemblages in Europe in response to climate change (Thuiller *et al.*, 2014). If the most sensitive butterfly species will go extinct, the associated loss of specific functional traits (or phylogenetic history) cannot be regained, which calls for careful conservation planning (Winter *et al.*, 2013).

Our results also highlight the potential cascading effect of plant invasions on multiple trophic levels. Indeed, invasive plant spread does not only lead to a decline in the phylogenetic diversity of plants (e.g. Winter *et al.*, 2009), but also to that of butterfly communities. This suggests that other herbivorous insect groups, such as beetles or flies, are likely affected similarly by invasive plants, which calls for repeated analyses of diverse insect

groups and potential effects on higher trophic levels (Bezemer *et al.*, 2014; Litt *et al.*, 2014; van Hengstum *et al.*, 2014).

To protect the Swiss butterfly diversity it is important to both prevent the introduction and spread of invasive plants and to protect the areas most at risk for dispersal-limited butterflies. Targeting areas for butterfly protection can be achieved by prioritizing sites (i) with the largest number of butterfly species identified as suffering from invasive plants (e.g. the Central Alps of Switzerland), and (ii) with the largest functional and phylogenetic diversities (e.g. using reserve selection algorithms; Arponen *et al.*, 2005; but see Winter *et al.*, 2013).

ACKNOWLEDGEMENT

We would like to thank Elena Concepción, Simona Gradinaru and Michael Nobis for helpful discussions on butterfly diversity, as well as BAFU, Hintermann and Weber for granting access to the BDM data and for extracting and formatting the data from the archive, respectively. We would also like to thank greatly the editor and two anonymous reviewers for their helpful and supportive comments. L.G. acknowledges support from the FORREG project (FOEN/WSL), N.E.Z. acknowledges support from the Swiss National Science Foundation (grant 31003A_149508/1). Funding to F.A. is from the Swiss National Science Foundation Grant PP00P3_150698.

SUPPORTING INFORMATION

Supporting information S1. Methodological details.

Supporting information S2. Results details.

BIOSKETCH

Laure Gallien's research generally aims at better understanding invasive species drivers and impacts. She likes exploring invasion patterns from different viewpoints, including coexistence theory, network theory and historical biogeography. At the moment she is particularly interested in developing a framework that can help anticipating invasive species impacts across trophic levels from the analysis of their functional trait. Author contributions: The study was designed by LG, and additional analyses suggested by NEZ, OS, and FA. NEZ, FA and MW collected the environmental, trait and phylogenetic data respectively. LG performed all analyses and wrote the first draft of the manuscript, which all authors edited.

REFERENCES

- Altermatt, F. (2010) Tell me what you eat and I'll tell you when you fly: diet can predict phenological changes in response to climate change. *Ecology Letters*, **13**, 1475–1484.
- Altermatt, F., Birrer, S., Plattner, M., Ramseier, P. & Stalling, T. (2008) Erste Resultate zu den Tagfaltern im Biodiversitätsmonitoring Schweiz. *Entomo Helvetica*, **1**, 75–83.
- Altermatt, F. & Pearse, I.S. (2011) Similarity and specialization of the larval versus adult diet of European butterflies and moths. *The American Naturalist*, **178**, 372–82.
- Arponen, A., Heikkinen, R.K., Thomas, C.D. & Moilanen, A. (2005) The Value of biodiversity in reserve selection: representation, species weighting, and benefit functions. *Conservation Biology*, **19**, 2009–2014.
- Benjamini, Y. & Yekutieli, D. (2001) The control of the false discovery rate in multiple testing under dependency. *The Annals of Statistics*, **29**, 1165–1188.
- Bezemer, T.M., Harvey, J.A. & Cronin, J.T. (2014) Response of native insect communities to invasive plants. *Annual Review of Ecology Evolution and Systematics*, **59**, 119–41.

440 Bolker, B. (2014) Dealing with quasi-models in R. *Creative Commons*, 1–5.

441 Brook, B.W., Sodhi, N.S. & Bradshaw, C.J.A. (2008) Synergies among extinction drivers
442 under global change. *Trends in Ecology and Evolution*, **23**, 453–460.

443 Cabra Rivas, I., Saldaña, A., Castro-Diez, P. & Gallien, L. (2015) A multi-scale approach to
444 identify invasion drivers and invaders' future dynamics. *Biological Invasions*, **2**, 411–
445 426.

446 Chytry, M., Maskell, L.C., Pino, J., Pysek, P., Vila, M., Font, X. & Smart, S.M. (2008)
447 Habitat invasions by alien plants: a quantitative comparison among Mediterranean,
448 subcontinental and oceanic regions of Europe. *Journal of Applied Ecology*, **45**, 448–458.

449 Concepción, E.D., Moretti, M., Altermatt, F., Nobis, M.P. & Obrist, M.K. (2015) Impacts of
450 urbanisation on biodiversity: the role of species mobility, degree of specialisation and
451 spatial scale. *Oikos*, **124**, 1571–1582.

452 Concepción, E.D., Obrist, M.K., Moretti, M., Altermatt, F., Baur, B. & Nobis, M.P. (2016)
453 Impacts of urban sprawl on species richness of plants, butterflies, gastropods and birds:
454 not only built-up area matters. *Urban Ecosystems*, **19**, 225–242.

455 Davies, K.F., Margules, C.R. & Lawrence, J.F. (2004) A synergistic effect puts rare,
456 specialized species at greater risk of extinction. *Ecology*, **85**, 265–271.

457 Davis, S.L. & Cipollini, D. (2014) Do mothers always know best? Oviposition mistakes and
458 resulting larval failure of *Pieris virginiensis* on *Alliaria petiolata*, a novel, toxic host.
459 *Biological Invasions*, **16**, 1941–1950.

460 Dray, S. & Dufour, A.B. (2007) The ade4 Package: Implementing the Duality Diagram for
461 Ecologists. *Journal of Statistical Software*, **22**, 1–20.

462 Dudley, R. & Srygley, R. (1994) Flight physiology of neotropical butterflies: allometry of

463 airspeeds during natural free flight. *Journal of Experimental Biology*, **191**, 125–139.

464 Ebert, G. (1991) *Die Schmetterlinge Baden-Württembergs*, Ebert. (ed. by Ulmer) Stuttgart,

465 Germany.

466 Ekroos, J., Heliölä, J. & Kuussaari, M. (2010) Homogenization of lepidopteran communities

467 in intensively cultivated agricultural landscapes. *Journal of Applied Ecology*, **47**, 459–

468 467.

469 Eskildsen, A., Carvalheiro, L.G., Kissling, W.D., Biesmeijer, J.C., Schweiger, O. & Høye,

470 T.T. (2015) Ecological specialization matters: long-term trends in butterfly species

471 richness and assemblage composition depend on multiple functional traits. *Diversity and*

472 *Distributions*, **21**, 792–802.

473 Forister, M.L., McCall, A.C., Sanders, N.J., Fordyce, J.A., Thorne, J.H., O’Brien, J., Waetjen,

474 D.P. & Shapiro, A.M. (2010) Compounded effects of climate change and habitat

475 alteration shift patterns of butterfly diversity. *Proceedings of the National Academy of*

476 *Sciences*, **107**, 2088–2092.

477 Forister, M.L. & Wilson, J.S. (2013) The population ecology of novel plant-herbivore

478 interactions. *Oikos*, **122**, 657–666.

479 Foster, J.T. & Robinson, S.K. (2007) Introduced birds and the fate of Hawaiian rainforests.

480 *Conservation Biology*, **21**, 1248–1257.

481 Freeman, E.A. & Moisen, G. (2008) PresenceAbsence: An R Package for Presence-Absence

482 Model Analysis. *Journal of Statistical Software*, **23**, 1–31.

483 Gallien, L. & Carboni, M. (2016) The community ecology of invasive species: Where are we

484 and what’s next? *Ecography*, In press, doi: 10.1111/ecog.02446.

485 Gallien, L., Mazel, F., Lavergne, S., Renaud, J., Douzet, R. & Thuiller, W. (2015) Contrasting

the effects of environment, dispersal and biotic interactions to explain the distribution of
invasive plants in alpine communities. *Biological Invasions*, **17**, 1407–1423.

Goslee, S.C. & Urban, D.L. (2007) The ecodist package for dissimilarity-based analysis of
ecological data. *Journal of Statistical Software*, **22**, 1–19.

Graves, S.D., Graves, S.D., Shapiro, A.M. & Shapiro, A.M. (2003) Exotics as host plants of
the California butterfly fauna. *Biological Conservation*, **110**, 413–433.

van Hengstum, T., Hooftman, D.A.P., Oostermeijer, J.G.B. & Tienderen, P.H. Van (2014)
Impact of plant invasions on local arthropod communities: a meta-analysis. *Journal of
Ecology*, **102**, 4–11.

Hulme, P.E. (2009) Trade, transport and trouble: managing invasive species pathways in an
era of globalization. *Journal of Applied Ecology*, **46**, 10–18.

Ignace, D.D. & Chesson, P. (2014) Removing an invader: evidence for forces reassembling a
Chihuahuan Desert ecosystem. *Ecology*, **95**, 3203–3212.

Jahner, J.P., Bonilla, M.M., Badik, K.J., Shapiro, A.M. & Forister, M.L. (2011) Use of Exotic
Hosts By Lepidoptera: Widespread Species Colonize More Novel Hosts. *Evolution*, **65**,
2719–2724.

Litt, A.R., Cord, E.E., Fulbright, T.E. & Schuster, G.L. (2014) Effects of Invasive Plants on
Arthropods. *Conservation Biology*, **28**, 1532–1549.

Masters, J.A. & Emery, S.M. (2015) The showy invasive plant *Ranunculus ficaria* facilitates
pollinator activity, pollen deposition, but not always seed production for two native
spring ephemeral plants. *Biological Invasions*, **17**, 2329–2337.

Morales, C.L., Arbetman, M.P., Cameron, S.A. & Aizen, M.A. (2013) Rapid ecological
replacement of a native bumble bee by invasive species. *Frontiers in Ecology and the*

509 *Environment*, **11**, 529–534.

510 Morales, C.L. & Traveset, A. (2009) A meta-analysis of impacts of alien vs. native plants on
511 pollinator visitation and reproductive success of co-flowering native plants. *Ecology*
512 *Letters*, **12**, 716–728.

513 Münkemüller, T., Boucher, F.C., Thuiller, W. & Lavergne, S. (2015) Phylogenetic niche
514 conservatism - common pitfalls and ways forward. *Functional Ecology*, **29**, 627–639.

515 Ohlemüller, R., Anderson, B.J., Araujo, M.B., Butchart, S.H., Kudrna, O., Ridgely, R.S. &
516 Thomas, C.D. (2008) The coincidence of climatic and species rarity: high risk to small-
517 range species from climate change. *Biology Letters*, **4**, 568–572.

518 Pavoine, S., Baguette, M., Stevens, V.M., Leibold, M. a., Turlure, C. & Bonsall, M.B. (2014)
519 Life-history traits, but not phylogeny, drive compositional patterns in a butterfly
520 metacommunity. *Ecology*, **95**, 3304–3313.

521 Payne, J.L. & Finnegan, S. (2007) The effect of geographic range on extinction risk during
522 background and mass extinction. *Proceedings of the National Academy of Sciences of*
523 *the United States of America*, **104**, 10506–10511.

524 Pearse, I.S. & Altermatt, F. (2013) Predicting novel trophic interactions in a non-native world.
525 *Ecology Letters*, **16**, 1088–1094.

526 Pyšek, P., Jarosik, V. & Kucera, T. (2003) Inclusion of native and alien species in temperate
527 nature reserves: an historical study from Central Europe. *Conservation Biology*, **17**,
528 1414–1424.

529 Richardson, D.M., Pyšek, P., Rejmánek, M., Barbour, M.G., Panetta, F.D. & West, C.J.
530 (2000) Naturalization and invasion of alien plants: concepts and definitions. *Diversity*
531 *and Distributions*, **6**, 93–107.

532 Sax, D.F. & Gaines, S.D. (2008) Species invasions and extinction: the future of native
533 biodiversity on islands. *Proceedings of the National Academy of Sciences of the United*
534 *States of America*, **105**, 11490–11497.

535 Schweiger, O., Biesmeijer, J.C., Bommarco, R., Hickler, T., Hulme, P.E., Klotz, S., Kühn, I.,
536 Moora, M., Nielsen, A., Ohlemüller, R., Petanidou, T., Potts, S.G., Pyšek, P., Stout, J.C.,
537 Sykes, M.T., Tscheulin, T., Vilà, M., Walther, G.R., Westphal, C., Winter, M., Zobel, M.
538 & Settele, J. (2010) Multiple stressors on biotic interactions: How climate change and
539 alien species interact to affect pollination. *Biological Reviews*, **85**, 777–795.

540 Schweiger, O., Harpke, A., Wiemers, M. & Settele, J. (2014) CLIMBER: Climatic niche
541 characteristics of the butterflies in Europe. *ZooKeys*, **367**, 65–84.

542 Sekar, S. (2012) A meta-analysis of the traits affecting dispersal ability in butterflies: can
543 wingspan be used as a proxy? *Journal of Animal Ecology*, **81**, 174–184.

544 Strobl, C., Boulesteix, A.-L., Zeileis, A. & Hothorn, T. (2007) Bias in random forest variable
545 importance measures: illustrations, sources and a solution. *BMC bioinformatics*, **8**, 25.

546 Tallamy, D.W. & Shropshire, K.J. (2009) Ranking Lepidopteran Use of Native Versus
547 Introduced Plants. *Conservation Biology*, **23**, 941–947.

548 Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011) MEGA5:
549 molecular evolutionary genetics analysis using maximum likelihood, evolutionary
550 distance, and maximum parsimony methods. *Molecular biology and evolution*, **28**,
551 2731–2739.

552 Thuiller, W., Pironon, S., Psomas, A., Barbet-Massin, M., Jiguet, F., Lavergne, S., Pearman,
553 P.B., Renaud, J., Zupan, L. & Zimmermann, N.E. (2014) The European functional tree
554 of bird life in the face of global change. *Nature Communications*, **5**, 3118.

- 555 Turlure, C., Schtickzelle, N. & Baguette, M. (2010) Resource grain scales mobility and adult
556 morphology in butterflies. *Landscape Ecology*, **25**, 95–108.
- 557 Venables, W.N. & Ripley, B.D. (2002) *Modern Applied Statistics with S*, 4th Editio. (ed. by
558 Springer) New York.
- 559 Vilà, M., Espinar, J.L., Hejda, M., Hulme, P.E., Jarosik, V., Maron, J.L., Pergl, J., Schaffner,
560 U., Sun, Y. & Pysek, P. (2011) Ecological impacts of invasive alien plants : a meta-
561 analysis of their effects on species , communities and ecosystems. *Ecology Letters*, **14**,
562 702–708.
- 563 Vilcinskis, a, Stoecker, K., Schmidtberg, H., Rohrich, C.R. & Vogel, H. (2013) Invasive
564 harlequin ladybird carries biological weapons against native competitors. *Science*, **340**,
565 862–863.
- 566 Villéger, S., Blanchet, S., Beauchard, O., Oberdorff, T. & Brosse, S. (2015) From current
567 distinctiveness to future homogenization of the world's freshwater fish faunas. *Diversity
568 and Distributions*, **21**, 223–235.
- 569 Walther, G.-R., Gritti, E.S., Berger, S., Hickler, T., Tang, Z. & Sykes, M.T. (2007) Palms
570 tracking climate change. *Global Ecology and Biogeography*, **16**, 801–809.
- 571 Warren, M.S., Hill, J.K., Thomas, J.A., Asher, J., Fox, R., Huntley, B., Roy, D.B., Telfer,
572 M.G., Jeffcoate, S., Harding, P., Jeffcoate, G., Willis, S.G., Greatorex-Davies, J.N.,
573 Moss, D. & Thomas, C.D. (2001) Rapid responses of British butterflies to opposing
574 forces of climate and habitat change. *Nature*, **414**, 65–69.
- 575 Webb, C.O., Ackerly, D.D., McPeck, M.A. & Donoghue, M.J. (2002) Phylogenies and
576 community ecology. *Annual Review of Ecology and Systematics*, **33**, 475–505.
- 577 Weed, A.S., Ayres, M.P. & Hicke, J.A. (2013) Consequences of climate change for biotic

disturbances in North American forests. *Ecological Monographs*, **83**, 441–470.

Winter, M., Devictor, V. & Schweiger, O. (2013) Phylogenetic diversity and nature conservation: where are we? *Trends in ecology & evolution*, **28**, 199–204.

Winter, M., Schweiger, O., Klotz, S., Nentwig, W., Andriopoulos, P., Arianoutsou, M., Basnou, C., Delipetrou, P., Didziulis, V., Hejda, M., Hulme, P.E., Lambdon, P.W., Pergl, J., Pysek, P., Roy, D.B. & Kühn, I. (2009) Plant extinctions and introductions lead to phylogenetic and taxonomic homogenization of the European flora. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 21721–21725.

Figure 1: Model response curves representing the variables to which the different species groups respond to. Native plants are represented in black, exotic plants in red, and native butterflies in blue. To facilitate comparisons between species groups and because the three taxonomic groups have different levels of species richness, we rescaled their responses between 0 and 100 (i.e. 100 representing the maximum observed richness). The shaded areas around the curves represent the 95% confidence interval around the model fitted values. Interpretations for the six PCA axes are indicated on the bottom arrow of each panel (see Fig. S1.3 for more detail).


Figure 2. Response curves of the butterfly richness model when incorporating the number of invasive plant species as a covariate. Only variables having a significant interaction term with the invasive plant richness are shown. In the lower right panel, the variable importance of the butterfly richness model is presented (the number of invasive species is highlighted in black). Increasing values along the six PCA axes can be interpreted as follows: (Axis1) decreasing temperature, increasing forest cover, increasing distance to urban and agricultural area, (Axis 2) increasing habitat diversity and proportion of urban area, (Axis 3) increasing

wetland cover, (Axis 4) increasing habitat diversity and precipitation, (Axis 5) increasing number of frost days, and (Axis 6) increasing levels of solar radiation (see Fig. S1.3 for more detail).

Figure 3. Characteristics and phylogenetic position of the butterfly species detected as suffering from invasive plant richness (note that none of the species was detected as profiting from invasive plant richness). (A) Difference in wing-load ratios between butterflies detected as significantly affected and those unaffected by invasive plant richness. (B) Phylogenetic position of the butterfly species detected as suffering from invasive plant richness (in red), not significantly affected (in black) and species not tested because of too few occurrences (in grey) among all Swiss butterfly species.


Figure 4. Spatial distribution of the butterfly species detected as suffering from the local richness in invasive plant species. Map backgrounds represent the average predicted richness of suffering species (white-to-blue gradient), while black dots of different size show their observed local richness (panel a) and proportion among the local butterfly richness (panel b).

619 **Figure 1.**



620
621
622


623 **Figure 2.**



624
625



629 **Figure 4.**



630
631